NIGERIA

## FAST FACTS FROM THE 2015 NIGERIA **MALARIA INDICATOR SURVEY**

The 2015 Nigeria Malaria Indicator Survey (NMIS) provides population-based estimates on malaria indicators including malaria prevention, case management, and malaria and anaemia prevalence to inform strategic planning and programme evaluation.

INSECTICIDE-TREATED NET (ITN) USE

Nearly **half** of children under 5 (**44%**) and pregnant women age 15-49 (**49%**) slept under an ITN the night before the survey.

IPTp

37% of women took 2 + doses of SP and received at least 1 during an antenatal care visit.

Trends in Intermittent Preventive Treatment of Pregnant Women (IPTp)

Percent of women age 15-49 with a live birth in the two years before the survey who took 2+ or 3+ doses of SP and received at least one during an antenatal care visit (ANC)

IPTp2+: 5, 13, 15, 37
IPTp3+: 2, 5, 6, 19

| 2008 NDHS	2010 NMIS	2013 NDHS	2015 NMIS

MALARIA PREVALENCE

More than **one-quarter** (**27%**) of children under 5 tested positive for malaria by microscopy.

Prevalence is **higher among rural households** than urban households.

Trends in Malaria Prevalence

Percent of children 6-59 months who tested positive for malaria by microscopy

■ 2010 NMIS ■ 2015 NMIS

	Total	Urban	Rural
2010 NMIS	42	23	48
2015 NMIS	27	12	36

USAID FROM THE AMERICAN PEOPLE

The Global Fund

USAID FROM THE AMERICAN PEOPLE — CDC — U.S. President's Malaria Initiative

UKaid from the British people

World Health Organization