

Disruption in HIV, Hepatitis and STI services due to COVID-19

Global HIV, Hepatitis and STI Programmes World Health Organization

Presentation with latest results as of July 1, 2020



Global COVID–19 Status – Case Counts

Status as of 1 July 2020 (6:08pm CEST) :

- 163 939 daily new cases
- 10 357 662 cumulative cases



Source: WHO Coronavirus Disease (COVID-19) Dashboard



Global COVID–19 Status - Deaths



Source: WHO Coronavirus Disease (COVID-19) Dashboard



Variation in Regional Trends

- Daily new infections are still fluctuating in all regions
- Increasing trend in the African, South-East Asia, American and Eastern Mediterranean Regions in the past few weeks.
- Plateauing trend in the Western Pacific and European Region in the past few weeks. However, fluctuations remain, for example some recent increases in daily cases in the European region





Countries adjust intensity of response over time

- COVID19 Government Response Stringency Index (Oxford University)
 - Composite measure based on nine response indicators including school closures, workplace closures, and travel bans
 - Darker blue = more stringent response



Source: Oxford University. Coronavirus Government Response Tracker. <u>https://www.bsg.ox.ac.uk/research</u>



Methods

- The World Health Organisation undertook a survey of relevant policies, drug stocks, service disruptions to treatment, testing and prevention and country responses, to inform WHO action to countries
- The data was provided by WHO country and regional offices for 144 countries and triangulated with other sources of **policy** and **health service disruption** data
 - Global AIDS Monitoring Data on the adoption and implementation of major HIV policies
 - A comprehensive WHO survey on impact of the pandemic on 25 essential health services across life course

• Survey undertaken 4th May to 30th June 2020, and is updated regularly for country action

- Critically low stock of ARVs is defined as stock of 3 months or less.
- Disruption of ARV services is reported by countries as a % of PLHIV on ART affected by disruption now. Risk of ARV disruption is forward looking risk of ARV disruption, as reported by WHO regional and country offices
- Disruption of services was collected for Prevention, Testing and Treatment Services for HIV, Hepatitis B and C, and Sexually Transmitted Infections, and for 25 essential health services within WHO (Health Service Performance Assessment, HSA)
- Data is updated regularly to facilitate WHO action for continuity of essential HIV, Hepatitis and STI services.
 ART manufacturing data is collected by WHO SEARO office



Questionnaire completion (as of June 2020)

Varies between regions and indicators

- MMD ARV most complete (includes GAM contribution)
- Strong data on ART stock from some regions, but less from others
- Incomplete or no data for some focus countries:

Angola	Kenya
Cameroon	Lesotho
Côte d'Ivoire	Malaysia
DRC	Mexico
Ghana	United Republic of Tanzania
Haiti	

Number of countries with data for each indicator



Source: Top graph: GAM UNAIDS/WHO/UNICEF (n=143) and WHO HIV/HEP/STI COVID-19 Questionnaire (n=84) Bottom graph: WHO HIV/HEP/STI COVID-19 Questionnaire



Disruption in Hepatitis Services



- **Argentina:** Overall reduction in service provision across health sector has impacted HBV and HCV testing and treatment initiation. There is a stock out of DAAs, and procurement process initiated.
- **Thailand:** The key services among higher risk groups such as HCV, HBV screening test and IEC in the community cannot be carried out in COVID-19 context
- **Russian Federation:** Viral hepatitis in Russia is exclusively treated infectionists specialists who are all now on engaged in COVID-19 treatment, this affects the situation with diagnosis and treatment of viral hepatitis
- **Hepatitis B:** Argentina, Belize, Botswana, Brazil, China, Ecuador, Ethiopia, Fiji,Kiribati Philippines, Russian Federation, Thailand, Vanuatu
- Hepatitis C: Argentina, Belize, Botswana, China, Costa Rica, Ethiopia, Kiribati, Philippines, Russian Federation, Thailand, Indonesia, Myanmar,



Source: WHO HIV/HEP/STI COVID-19 Questionnaire

Disruption in STI services

 STI service disruption in 16 countries (including PMTCT for syphilis, possible drug stock outs, testing kits stock outs), of which:

Botswana	Ethiopia	Papua New Guinea
China	Guatemala	Philippines
Ecuador	Honduras	Uganda
El Salvador	Nigeria	

• Ecuador: Country has 1 month-stock of RDT for syphilis (it may impact EMTCT; but there are alternative testing available)

• Honduras: Stock-out of syphilis RDT; 2month-stock of VL; 3 months CD4; backload in genotyping and zero stock



ARV multi month dispensing

- ARV MMD policy is adopted in most countries.
- Data available for 144 countries:
 - 129 (90%) adopted MMD policy
- County cases suggest COVID-19 effect on MMD is double-edged:
 - Sufficient ARV stock → intensified MMD (Namibia, Malawi...)
 - Uncertain ARV stock → shorter MMD (Indonesia, Botswana..)





Countries with MMD policy per WHO region: frequency of ARV pick-up



Source: GAM UNAIDS/WHO/UNICEF and WHO HIV/HEP/STI COVID-19 Questionnaire



MMD for other HIV/Hepatitis/STI drugs



Source: WHO HIV/HEP/STI COVID-19 Questionnaire



Countries are balancing stock levels with move to MMD

• Eswatini

"There are ongoing in-country discussions on providing multi month (3-6 months) scripts for ART/TB/NCD patients; the central medical store is yet to advise if the current stock levels can allow for this initiatives."

United Republic of Tanzania:

"3 monthly dispensing country widespread. 6 months have been started in Dar Es Salaam but could revert to 3 month due to stock shortages"

South Africa:

"(ARV Picky up frequency of) Every 3 months, Western Cape is giving 4 months, 2 months TEE (tenofovir-emtricitabine-efavirenz) due to low stock."



ARV stock availability of first line stocks

- Data available for 84 countries
- 24 countries reported ARV stocks availability for major first line drugs (TLE/TEE/TLD) of **3 months or less**
 - 13 (54%) at medium/high risk of stock outs:



Source: WHO HIV/HEP/STI COVID-19 Questionnaire



Other possible stock-outs

• Possible ARV 2nd line stock-outs in 27 countries:

LPV/r :	Bhutan, Botswana, Cuba, Ecuador, El Salvador, Honduras, Kenya, Myanmar, Nepal, Paraguay, Russian Federation, Sri Lanka, Suriname, Timor Leste (14)
ATV/r :	Cuba, El Salvador, Jamaica, Kenya, Peru, Venezuela, Zambia (7)
Pediatric ARV:	Botswana, Cuba, Ecuador, El Salvador, Honduras, Jamaica, Namibia, Nepal, Nicaragua, Paraguay, Panama, Peru, Suriname, Zimbabwe (14)

- Other ARV drugs with possible stock outs: **AZT/3TC, Abacavir, DRV** in some countries.
- Hepatitis C Drugs : Argentina, Indonesia
- STI Treatment : Madagascar, Papua New Guinea, Uganda
- OI drugs : Eswatini, Kenya
- Syphilis test kits : Ethiopia, Uganda



Monitoring risk of ARV disruptions

 73 countries reported risk of ARV disruptions – in varying degrees (41% medium/high risk.)*

Cited causes:

- "Failure of suppliers to deliver on time"
- "Courier services shut down due to COVID"
- "Government has to pay upfront payments in order for some supplies to process order."





Covid-19 and Impact on HIV drug manufacturers

Scenario in India: Mar - Apr 2020

- Initial <u>ban by Government</u> barring export of drugs including ARVs; but lifted after 2 weeks
- Discussions in March 2020 with manufacturers:
- Adequate <u>API stocks</u> to meet demand.
- Other issues highlighted were:
 - Reduced workforce in plants, and to load to trucks;
 - <u>Ground Transportation</u> issues to reach warehouses and beyond, despite Government clarification allowing transport of essential goods;
 - Shortage of <u>packaging</u> material; and
 - Reduced movement of <u>cargo flights</u>.
- Successful facilitation in shipping drugs to Nepal by road and Indonesia by sea route to address impending stockouts.

Scenario in May 2020

- Most of road transport is normal now
- Many cargo flights
- No ban on exports of ARVs
- GF Assessment Moderate Impact overall (details summarized in separate slide)

Current Scenario – Detailed Assessment

- Email-based interaction with leading manufacturers
- Structured exchange on issues faced
- Feedback on challenges as well as suggestions
- 4/9 responses in 2 days; others have committed too
- Details of initial responses summarized in next slide



Countries reporting on ARV disruptions due to COVID-19, 2020

Preliminary results compiled from surveys conducted by WHO between April and June 2020 (n=127)



Disclaimer: The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.



PLHIV on ART affected due to COVID19

Now

- 36 countries reported disruption in provision of ARV services. In these countries 11.5 million people were receiving ART (45% of total PLHIV on ART))
- Ranges from <1% 40% of PLHIV on ART in a country
- 24 countries had a critically low stock of ARVs. In these countries 8.3 million people were receiving ART (33% of total PLHIV on ART)

In 3 – 6 months

 73 countries are facing risk of ARV disruption. In these countries 17.7 million people were receiving ART (70% of total PLHIV on ART)



Disruptions in other services due to COVID-19



Prevention programs for VMMC and PWID are in selected countries, so disruption may be in most countries where there is a program

Source: WHO HIV/HEP/STI COVID-19 Questionnaire



- "Regular communications from Ministry of health HIV department in the form of circular letters to all service providers, toll free phone lines for consultations and support, flexed criteria for six months dispensation, running HIV clinics every day to reduce congestion, suspension of non essential HIV services that need frequent visits to the facility. – Malawi
- **"Implemented intensified MMD** for both adults and paediatrics. Decongesting ART facilities, Establishment of a **hotline for PLWHIV**, community dispensing at existing DSD model sites and primary health care outreach points, Hiring of short term staff to ensure service continuity." **Namibia**
- "Essential Service Advisory A separate interim guidance on continuity of ART and OST was issued by the Ministry. ART delivered by road not plane in May from Mumbai. OST takeaway dose for 90% of clients with careful monitoring and contact by phone with weekly updates" **Nepal**
- "Due to lockdown, limited access of PLHIV to treatment centers. NAP ensuring **home delivery of ARVs** via courier." **Pakistan**
- "National HIV program is planning to **address on HIV viral load testing by using Xpert**, that are already operating in every provinces, to mitigate delay and disruption of sample transportation due to COVID 19." Lao People's Democratic Republic
- "Strengthened out outreach interventions through peer and mobile outreach, PrEP is ordered online and delivered by mobile team also." - Georgia



Conclusions

HIV, Hepatitis and STI service COVID disruptions due to

- 1. Stock out of drugs
- 2. Diversion of health systems capacity
- 3. Lack of access to health facilities during lock down.

Widespread ARV disruptions have been reported to WHO across all regions

- 73 countries are facing risk of ARV disruption. In these countries 17.7 million people were receiving ART (70% of total PLHIV on ART)
- 36 countries reported disruption in provision of ARV services. In these countries 11.5 million people were receiving ART (45% of total PLHIV on ART))
- 24 countries had a critically low stock of ARVs. In these countries 8.3 million people were receiving ART (33% of total PLHIV on ART)

Additional disruptions on HIV testing, prevention, Hepatitis and STI services

- 61 countries facing disruption in other prevention, testing and treatment services
- 38 countries report disruption in HIV testing, 17 in Key Population services, 12 in condom provision, and 7 in PrEP
- 10 countries report disruption in Hepatitis B testing, 6 in Hepatitis C Treatment initiation, and 5 in Family Planning

• Country responses have mitigated the worst case service disruptions but risks remain widespread

- 90% of countries adopted policy of multi month dispensing of ARVs
- Countries are balancing stock levels with a shift to MMD, with some countries moving to 6 months and then reverting to 3 months due to stock shortages
- Continuity of essential package of health services requires policy and financial support for key services
- Innovative Service Delivery use of community delivery, MMD for IDU and Hepatitis C treatment, may have benefits in the long run by reducing clinical staff times and promoting innovation in people centred service delivery

